



p-ISSN 0854-1418

e-ISSN 2503-426X

Accreditation by the Directorate General of Higher Education,  
Research and Technology No. 158/E/KPT/2021

# EKSPLORIUM

**Buletin Pusat Pengembangan Bahan Galian Nuklir**

Volume 43 No. 1, May 2022



**DIRECTORATE OF REPOSITORIES, MULTIMEDIA, AND SCIENTIFIC PUBLICATION  
NATIONAL RESEARCH AND INNOVATION AGENCY**

Eksplorium	Volume 43	No. 1	Pages 1-58	Jakarta May 2022	p-ISSN 0854- 1418 e-ISSN 2503-426X	Accreditation No. 158/E/KPT/2021
------------	--------------	----------	---------------	---------------------	---------------------------------------	-------------------------------------



9 772503 426021



9 770854 141426

**p-ISSN 0854-1418**  
**e-ISSN 2503-426X**

Accreditation by the Directorate General of Higher Education,  
Research and Technology No. 158/E/KPT/2021

# EKSPLORIUM

---

## Buletin Pusat Pengembangan Bahan Galian Nuklir

---

**Volume 43 No. 1, May 2022**

**Eksplorium** is Buletin Pusat Pengembangan Bahan Galian Nuklir as information media which contains the results of studies, research, and development of nuclear geology with the scope of geology, exploration, mining, processing of nuclear minerals, and environmental safety as well as the development of nuclear technology for people's welfare  
**Eksplorium** published 2 (two) times a year, in May and November.

### **Chief Editor**

Frederikus Dian Indrastomo, M.T., BRIN, Indonesia

### **Associate Editors**

I Gde Sukadana, M.Eng., BRIN, Indonesia  
Heri Syaeful, M.T., BRIN, Indonesia  
Kurnia Setiawan Widana, M.T., BRIN, Indonesia  
Dwi Haryanto, M.Si., BRIN, Indonesia  
Mirna Berliana Garwan S.T., BRIN, Indonesia  
Windi Anarta Draniswari, S.T., BRIN, Indonesia  
Ersina Rakhma, M.T., BRIN, Indonesia  
Wira Cakrabuana, S.T., BRIN, Indonesia

### **Editorial Board**

Prof. Dr. Muhayatun Santoso, M.T., BRIN, Indonesia  
Dr. Eng. Imam Achmad Sadisun, Bandung Institute of Technology (ITB), Indonesia  
Dr. I Wayan Warmada, Universitas Gadjah Mada (UGM), Indonesia  
Dr. Sri Mulyaningsih, IST "AKPRIND" Yogyakarta, Indonesia  
Dr. Hill Gendoet Hartono, Yogyakarta National Institute of Technology (ITNY), Indonesia  
Dr. Sulaksana Permana, Universitas Indonesia (Alumni), Indonesia  
Iwan Susanto, Ph.D., Jakarta State Polytechnic (PNJ), Indonesia  
Yuniarti Ulfa, M.Sc., Polytechnic of Geology and Mining (AGP), Indonesia

Photo caption: Tanjunggenting Reverse Fault, Bangka (Contributor:Heri Syaeful)

---

### **Publisher:**

**DIRECTORATE OF REPOSITORIES, MULTIMEDIA, AND SCIENTIFIC PUBLICATION**  
**NATIONAL RESEARCH AND INNOVATION AGENCY**

---

Editorial Address:

Building 20, B.J. Habibie Science and Technology Area, South Tangerang, Banten 15314

Telp. 081 110 646 830, E-mail: [eksplorium@brin.go.id](mailto:eksplorium@brin.go.id)

Website: <http://jurnal.batan.go.id/index.php/eksplorium>

---

# EKSPLORIUM

Buletin Pusat Pengembangan Bahan Galian Nuklir  
Volume 43, No. 1, May 2022

---



---

## Table of Content

Foreword .....	i
Table of Content.....	ii
Current Content.....	iii
Review on Granitic Rocks in Sumatra: Intrusion Process, Classification, Mineralization, and Potential Uses. <i>Ronaldo Irzon, Heri Syaeful, Aries Kusworo, Joko Wahyudiono, Ngadenin</i> .....	1–12
Ore Mineralization Characteristics in Hydrothermal Alteration at Mangunharjo and Surrounding Area, Pacitan, Indonesia. <i>Abdul Faisal Baba, Sri Mulyaningsih, Radhitya Adzan Hidayah</i> .....	13–22
Sub-surface Geological Modeling Based on Gravity Residual Data in Adang Volcanic Rock Area, Mamuju, West Sulawesi Province. <i>Adhika Junara Karunianto, Dwi Haryanto, Ngadenin</i> .....	23–28
Tectonic Pattern Imaging of Southern Sumatra Region Using Double Difference Seismic Tomography. <i>Akmal Firmansyah, Wandono, Mohamad Ramdhan</i> .....	29–40
Geostatistical Ore Body Modeling on Uranium Mineralization in Remaja Sector, Kalan Area, West Kalimantan. <i>Roni Cahya Ciputra, Mohamad Nur Heriawan, Heri Syaeful, Dhatu Kamajati, Putri Rahmawati</i> .....	41–58

# EKSPLORIUM

Buletin Pusat Pengembangan Bahan Galian Nuklir  
Volume 43, No. 1, May 2022

---

---

## Review on Granitic Rocks in Sumatra: Intrusion Process, Classification, Mineralization, and Potential Uses

Ronaldo Irzon<sup>1\*</sup>, Heri Syaeful<sup>2</sup>, Aries Kusworo<sup>1</sup>, Joko Wahyudiono<sup>1</sup>, Ngadenin<sup>2</sup>

<sup>1</sup>Center for Geological Survey, Geological Agency, Ministry of Energy and Mineral Resources  
Diponegoro St., No. 57, Bandung 40122, Indonesia

<sup>2</sup>Research Center for Nuclear Fuel Cycle and Radioactive Waste Technology,  
Research Organization for Nuclear Energy (ORTN), National Research and Innovation Agency (BRIN)  
PUSPIPTEK Area Building 20, South Tangerang 15314, Indonesia

\*E-mail: [ronaldoirzon18@gmail.com](mailto:ronaldoirzon18@gmail.com)

Article received: 20 September 2021, revised: 22 April 2022, accepted: 28 April 2022

DOI: 10.17146/eksplorium.2022.43.1.6471

### ABSTRACT

Granitic rocks are widely distributed in Sumatra and surrounding areas. These granitoids are classified into several granite provinces of Southeast Asia with different intrusion processes and specific characteristics. This paper aims to review the intrusion of granitic rocks in Sumatra and describe the opportunities associated with it. Granite rocks are used to manufacture cultural heritage, works of art, and ornaments because of their weathering resistance, color diversity, and hardness characters. S-type granite intrusion in Sumatra might be associated with tin mineralization while silver-gold with the I-type. Theoretically, granite contains more REE than other igneous rocks. Mining and extraction difficulties complicate the direct REE exploitation from fresh granite. A-type granite relatively contains more REE than the other types, but this type of granite is not correlated with certain provinces. Indonesia has a tropical climate which is prone to weathering. Therefore, it is possible for REE and/or bauxite enrichments in the granite weathering horizon. Granite is assumed to be a potential source of uranium and thorium, especially for the S-type, because it is formed through the compression of sediments that can absorb these radioactive elements from the continental crust.

**Keywords:** Sumatra, granite, classification, mineralization

---

# EKSPLORIUM

Buletin Pusat Pengembangan Bahan Galian Nuklir  
Volume 43, No. 1, May 2022

---

---

## Ore Mineralization Characteristics in Hydrothermal Alteration at Mangunharjo and Surrounding Area, Pacitan, Indonesia

**Abdul Faisal Baba\***, Sri Mulyaningsih, Radhitya Adzan Hidayah

Geological Engineering IST AKPRIND, Yogyakarta  
I Nyoman Oka St. No.32, Yogyakarta, 55224, Indonesia

E-mail: [abdulfaisal13497@gmail.com](mailto:abdulfaisal13497@gmail.com)

Article received: 13 January 2021, revised: 2 Mei 2021, accepted: 28 March 2022

DOI: 10.17146/eksplorium.2022.43.1.6194

### ABSTRACT

The research area is located in Mangunharjo-Grindulu, Pacitan (Indonesia), as part of the Southern Mountain Tertiary Volcanic Arch. Outcrops of quartz veins-riched volcanic rock associated with sulfide minerals are found in this area. The Southern Mountain Oligo-Miocene magmatic arc is known as the potential area that contains precious metal deposits. The study aimed to determine the characteristics of the mineralized zone in this area. The research methods are geological surface mapping, thin-section observation, mineragraphy, and X-Ray Diffraction (XRD). The results show that the constituent lithologies were andesitic lava, breccia, and tuff; co-ignimbrite breccia, dacitic pumice and tuff, and dacitic dike; and pyroxene-rich andesitic volcanic rocks. The geological structure is dominated by oblique normal faults, strike-slip faults, and upward oblique faults associated with shear joints filled with quartz veins. Fieldwork observation, thin-section analyses, and mineragraphic and XRD observations identify three alteration zones in the hydrothermal system: the advanced argillic zone, the intermediate argillic zone, and the chloritized zone. By the mineral's association, it is interpreted that the advanced argillic zone was formed at a temperature of 220-330°C and pH 3-6 due to dissemination with side rocks located near the hydrothermal flows; the intermediate argillic zone and the chloritized zone were formed at a temperature of 150-300°C and a pH of 5-6 due to chloritized alteration of the hydrothermal fluid carrying the ore. This alteration zone has no economic potential for precious metal minerals so it is better to be developed for education, conservation, and natural laboratories.

**Keywords:** Geology, Ore Mineralization, Hydrothermal Alteration

---

# EKSPLORIUM

Buletin Pusat Pengembangan Bahan Galian Nuklir  
Volume 43, No. 1, May 2022

---

---

## Sub-surface Geological Modeling Based on Gravity Residual Data in Adang Volcanic Rock Area, Mamuju, West Sulawesi Province

Adhika Junara Karunianto<sup>1\*</sup>, Dwi Haryanto<sup>2</sup>, Ngadenin<sup>3</sup>

<sup>1</sup>Research Center for Mining Technology

Research Center for Nano Material Technology, National Research and Innovation Agency (BRIN)  
Building 820, PUSPIPTEK Area, Tangerang Selatan, Banten 15314, Indonesia

<sup>2</sup>Directorate of Laboratory Management, Research Facilities, and Science and Technology Park  
National Research and Innovation Agency (BRIN)

Grafika Sekip St., No. 2, Sleman, D. I. Yogyakarta 55284, Indonesia

<sup>3</sup>Research Center for Nuclear Fuel Cycle and Radioactive Waste Technology  
Research Organization for Nuclear Energy (ORTN), National Research and Innovation Agency (BRIN)

Building 20, PUSPIPTEK Area, Tangerang Selatan, Banten 15314, Indonesia

\*E-mail: [adhi007@brin.go.id](mailto:adhi007@brin.go.id)

Accepted: 3 November 2020, revised: 28 November 2021, accepted: 28 April 2021

DOI: 10.17146/eksplorium.2022.43.1.6089

### ABSTRACT

The Mamuju area of West Sulawesi Province is composed of Adang volcanic rock that is a product of the process of volcanism in a volcanic complex with an eruption center and several lava domes. The geology of the study area is composed of eleven rock units, namely Adang breccia, Adang lava, lava dome, volcanic conglomerate, Ampalas breccia, Malunda breccia, Boteng lava, Tapalang breccia, limestone, reef limestone, and alluvium. The mineralization of uranium, thorium, and rare earth elements formed in Adang lava thorianite veins. Adang lava is intruded by a dioritoid found in the Mamuju river upstream. The gravity modeling technique has produced two 2-D subsurface models based on gravity data on 2-D cross-sections of the residual gravity map. It is known that the rock density range from 2.10 to 2.85 g/cm<sup>3</sup> in the study area. Based on the interpretation of two 2-D subsurface models, a batholith, a giant-sized intrusive rock, is found in the southeastern part of the study area with a rock density of about 2.85 g/cm<sup>3</sup> and is interpreted to be dioritoids. Furthermore, deep intrusion rocks also occur in the center part of the study area with a rock density of 2.8 g/cm<sup>3</sup>. It is estimated to be the same as the rock in the southeastern part of the study area. Dioritoid intrudes a volcanic breccia sedimentary rock with a density of about 2.1 g/cm<sup>3</sup>.

**Keywords:** Mamuju, gravity, modeling, density, batholith

---

# EKSPLORIUM

Buletin Pusat Pengembangan Bahan Galian Nuklir  
Volume 43, No. 1, May 2022

---

---

## Tectonic Pattern Imaging of Southern Sumatra Region Using Double Difference Seismic Tomography

Akmal Firmansyah<sup>1</sup>, Wandono<sup>2</sup>, Mohamad Ramdhan<sup>3\*</sup>

<sup>1</sup>Center for Earthquake and Tsunami

Indonesian Agency for Meteorology, Climatology, and Geophysics (BMKG),  
Angkasa I St., No. 2, Kemayoran, Jakarta 10720, Indonesia

<sup>2</sup>Study Program of Geophysics; School of Meteorology, Climatology, and Geophysics of Indonesia (STMKG),  
Pondok Betung, Pondok Aren, Tangerang Selatan, Banten 15221, Indonesia

<sup>3</sup>Research Center for Geological Disaster, National Research and Innovation Agency (BRIN),  
Sangkuriang St., BRIN Science and Technology Area, Bandung, 40135, Indonesia

\*E-mail: [mohamad.ramdhan@brin.go.id](mailto:mohamad.ramdhan@brin.go.id)

Article received: 15 March 2022, revised: 12 Mei 2022, accepted: 31 Mei 2022

DOI: 10.17146/eksplorium.2022.43.1.6603

### ABSTRACT

Southern Sumatra and its surroundings are close to the contact zone of the Indo-Australian plate and Eurasian plate, so the area always relates to the high seismicity zone. The Sumatran subduction zone, the Mentawai fault, and several segments of the Sumatran fault drive seismic activities in the area. Tectonic settings are essential to understanding the area's source and hazard. This understanding can be obtained using the relocated hypocenter distribution and the 3D velocity model in the area. Relocated hypocenters and velocity models are obtained from simultaneous inversion from the BMKG earthquake catalog in January 2012-December 2020 using the double difference seismic tomography method. Seismic velocity inversion of P- and S- wave tomograms image the thermal zone beneath Dempo and Patah volcanoes at a depth of 30-50 km. Slab dehydration is also observed in several forearc high zone. Both phenomena are associated with negative anomalies. The Sumatran and Mentawai fault zones are marked between negative and positive anomalies on the contact zone. The subducted slab of the Indo-Australian plate is observed until a depth of 150 km, which is the maximum depth of nodes used in this study. The granitic basement beneath Anak Krakatau volcano is detected until 10 km. Two of those geological features are related to positive anomalies.

**Keywords:** seismic tomography, double difference, Southern Sumatra

---

# EKSPLORIUM

Buletin Pusat Pengembangan Bahan Galian Nuklir  
Volume 43, No. 1, May 2022

---

---

## Geostatistical Ore Body Modeling on Uranium Mineralization in Remaja Sector, Kalan Area, West Kalimantan

Roni Cahya Ciputra<sup>1,2\*</sup>, Mohamad Nur Heriawan<sup>1</sup>, Heri Syaeful<sup>2</sup>, Dhatu Kamajati<sup>3</sup>, Putri Rahmawati<sup>3</sup>

<sup>1</sup>Faculty of Mining and Petroleum Engineering, Bandung Institute of Technology (ITB)  
Ganesha St., No. 10, Bandung 40132, Indonesia

<sup>2</sup>Research Center for Nuclear Fuel Cycle and Radioactive Waste Technology,  
Research Organization for Nuclear Energy (ORTN), National Research and Innovation Agency (BRIN)  
PUSPIPTEK Area, Building 20, South Tangerang 15314, Indonesia

<sup>3</sup>Bureau for Public Communication, General Affairs, and Secretariat,  
National Research and Innovation Agency (BRIN)  
B.J. Habibie Building, M.H. Thamrin St., No. 8, Central Jakarta 10340, Indonesia

\*E-mail: [roni.cahya.ciputra@brin.go.id](mailto:roni.cahya.ciputra@brin.go.id)

Article received: 18 May 2022, revised: 24 May 2022, accepted: 31 May 2022

DOI: 10.17146/eksplorium.2022.43.1.6622

### ABSTRACT

Manual ore body modeling on Remaja Sector, Kalan, West Kalimantan generally takes a long time and is subjective. On the other hand, automatic modeling (implicit modeling) is faster, objective, and equipped with uncertainty factors. This study aimed to analyze the comparison between the geostatistical Sequential Indicator Simulation (SIS) ore body model to the manual ore body model. The lithology database was used as input for variogram analysis and SIS simulation. The directional variogram was used to construct an experimental variogram for the lithology with orientation data. The orientation of the lithologies corresponds to the anisotropy of their variogram map. The SIS was carried out in Block A and Block B with block sizes of  $6 \times 6 \times 6 \text{ m}^3$  and  $5 \times 5 \times 5 \text{ m}^3$  respectively. The simulation results were processed to produce a lithology probability model. By using maximum probability as block lithology, simulation results were well validated by the composite database histogram, the lithologies along the tunnel on the geological map of level 450 masl of Eko Remaja Tunnel., and the lithologies along boreholes. The weakness of the geostatistical ore body model was the results depending on the input parameters. Meanwhile, several advantages of the geostatistical ore body model were a faster processing process, equipped with an uncertainty factor, and the block size of the model has taken into account the distance between grade data so that it can be used directly for grade estimation. Quantitatively, the geostatistical ore body model had a higher average percentage of conformity to the lithology of the mineralized zone along the borehole than the manual ore body model.

**Keywords:** Kalan, geological modeling, geostatistics, sequential indicator simulation, uranium



# EKSPLORIUM

## **Buletin Pusat Pengembangan Bahan Galian Nuklir**

### MANUSCRIPT WRITING GUIDELINES

The articles published in this journal are the results of studies, research, and development of nuclear geology with the scope of geology, exploration, mining, processing of nuclear minerals, and environmental safety as well as the development of nuclear technology for people's welfare. The articles are original work and have never been published.

Article formats:

1. TITLE, capitalized in the beginning of each word with font size of 12 in the center position.
2. AUTHOR NAME, written 2 space below the title with font size of 10
3. AFFILIATION ADDRESS AND CORRESPONDING E-MAIL ADDRESS, written a space below the author name with font size of 10
4. ABSTRACT, written in English with a maximum of 200 words containing a summary of: problems, objectives, methods, results, and conclusions, also equipped with 3-5 keywords .
5. INTRODUCTION, contains the background, scope, and objectives of the research.
6. THEORY, if necessary.
7. A. METHODOLOGY for research scientific work;  
B. MAIN SUBJECT for review scientific work.
8. RESULT AND DISCUSSION, the results are arranged in detail containing data in the form of tables and figures, while the discussion of the results obtained is discussed based on basic concepts or hypotheses.
9. CONCLUSION, contains conclusions from research results and suggestions.
10. BIBLIOGRAPHY, written in the referring order and using serial numbers with numbering according to the rules. Example:
  - [1] A. El Taher, "Elemental Analysis of Granite by Instrumental Neutron Activation Analysis (INAA) and X-Ray Fluorescence Analysis (XRF)", *Appl.Radiat.Isot*, vol.70, p.350-354, 2012.
  - [2] F. Ferrari, T. Apuani, and G.P. Giani, "Rock Mass Rating Spatial Estimation by Geostatistical Analysis", *Int. J. Rock Mec. Min. Sci*, vol. 70, p. 162-176, 2014.
  - [3] L. Blevin, "Metallogeny of Granitic Rocks", *The Ishihara Symposium: Granites and Associated Metallogenesis*, Geoscience Australia, p. 1-4, 2004.
  - [4] H. Syaeful, Suharji, and A. Sumaryanto, "Pemodelan Geologi dan Estimasi Kalan, Kalimantan Barat", *Prosiding Seminar Nasional Teknologi Energi Nuklir*, Pontianak, 2014.

The bibliography contains at least 10 references for research scientific work and a minimum of 25 references for review scientific work, published in the last 10 years and at least 80% comes from primary references (journals and theses).

The articles are written in English on A4 paper with a maximum number of 15 pages including pictures and tables. The complete templates and writing instructions can be downloaded on the explorium page (<http://jurnal.batan.go.id/index.php/eksplorium>). The articles are submitted to the Editor in soft copy format on the Open Journal System/OJS of Eksplorium page no later than two months before the scheduled publication. Complete information can be read on the explorium page.